

Mass Generation and Non Commutative Spacetime

B.G. Sidharth

International Institute for Applicable Mathematics & Information Sciences
Hyderabad (India) & Udine (Italy)

B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

Abstract

In this paper we show how it is possible to obtain mass generation in the context of non Abelian gauge field theory, using a non commutative spacetime. This is further confirmed by the modified dispersion relation that results from such a geometry.

1 Introduction

It is well known that Hermann Weyl's original phase transformation proposal was generalized, so that the global or constant phase of λ was considered to be a function of the coordinates [1, 2, 3, 4].

As is well known this leads to a covariant gauge derivative. For example, the transformation arising from $(x^\mu) \rightarrow (x^\mu + dx^\mu)$,

$$\psi \rightarrow \psi e^{-i\lambda} \quad (1)$$

leads to the familiar electromagnetic potential gauge,

$$A_\mu \rightarrow A_\mu - \partial_\mu \lambda \quad (2)$$

The above transformation, ofcourse, is a symmetry transformation. In the transition from (1) to (2), we expand the exponential, retaining terms only to the first order in coordinate differentials.

Let us now consider the gauge field in some detail. As is known this could be

obtained as a generalization of the above phase function λ to include fields with internal degrees of freedom. For example λ could be replaced by A_μ given by [2]

$$A_\mu = \sum_i A_\mu^i(x) L_i, \quad (3)$$

The gauge field itself would be obtained by using Stoke's Theorem and (3). This is a very well known procedure: considering a circuit, which for simplicity we can take to be a parallelogram of side dx and dy in two dimensions, we can easily deduce the equation for the field, viz.,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - iq[A_\mu, A_\nu], \quad (4)$$

q being the gauge field coupling constant.

In (4), the second term on the right side is typical of a non Abelian gauge field. In the case of the U(1) electromagnetic field, this latter term vanishes. Further as is well known, in a typical Lagrangian like

$$L = \bar{\psi} \gamma^\mu D_\mu \psi - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} - m \bar{\psi} \psi \quad (5)$$

D denoting the Gauge covariant derivative, there is no mass term for the field Bosons. Such a mass term in (5) must have the form $m^2 A^\mu A_\mu$ which unfortunately is not Gauge invariant.

This was the shortcoming of the original Yang-Mills Gauge Theory: The Gauge Bosons would be massless and hence the need for a symmetry breaking, mass generating mechanism.

The well known remedy for the above situation has been to consider, in analogy with superconductivity theory, an extra phase of a self coherent system (Cf.ref.[2] for a simple and elegant treatment and also refs. [3] and [5]). Thus instead of the gauge field A_μ , we consider a new phase adjusted gauge field after the symmetry is broken

$$W_\mu = A_\mu - \frac{1}{q} \partial_\mu \phi \quad (6)$$

The field W_μ now generates the mass in a self consistent manner via a Higgs mechanism. Infact the kinetic energy term

$$\frac{1}{2} |D_\mu \phi|^2 \quad , \quad (7)$$

where D_μ in (7) denotes the Gauge , now becomes

$$|D_\mu \phi_0|^2 = q^2 |W_\mu|^2 |\phi_0|^2, \quad (8)$$

Equation (8) gives the mass in terms of the ground state ϕ_0 .

The whole point is as follows: The symmetry breaking of the gauge field manifests itself only at short length scales signifying the fact that the field is mediated by particles with large mass. Further the internal symmetry space of the gauge field is broken by an external constraint: the wave function has an intrinsic relative phase factor which is a different function of spacetime coordinates compared to the phase change necessitated by the minimum coupling requirement for a free particle with the gauge potential. This cannot be achieved for an ordinary point like particle, but a new type of a physical system, like the self coherent system of superconductivity theory now interacts with the gauge field. The second or extra term in (6) is effectively an external field, though (8) manifests itself only in a relatively small spatial interval. The ϕ of the Higgs field in (6), in analogy with the phase function of Cooper pairs of superconductivity theory comes with a Landau-Ginzburg potential $V(\phi)$.

Let us now consider in the gauge field transformation, an additional phase term, $f(x)$, this being a scalar. In the usual theory such a term can always be gauged away in the U(1) electromagnetic group. However we now consider the new situation of a noncommutative geometry viz.,

$$[dx^\mu, dx^\nu] = \Theta^{\mu\nu} \beta, \beta \sim 0(l^2) \quad (9)$$

where l denotes a minimum spacetime cut off. Equation (9) is infact Lorentz covariant. Then the f phase factor gives a contribution to the second order in coordinate differentials,

$$\begin{aligned} & \frac{1}{2} [\partial_\mu B_\nu - \partial_\nu B_\mu] [dx^\mu, dx^\nu] \\ & + \frac{1}{2} [\partial_\mu B_\nu + \partial_\nu B_\mu] [dx^\mu dx^\nu + dx^\nu dx^\mu] \end{aligned} \quad (10)$$

where $B_\mu \equiv \partial_\mu f$.

As can be seen from (10) and (9), the new contribution is in the term which contains the commutator of the coordinate differentials, and not in the symmetric second term. Effectively, remembering that B_μ arises from the scalar

phase factor, and not from the non-Abelian gauge field, in equation (4) A_μ is replaced by

$$A_\mu \rightarrow A_\mu + B_\mu = A_\mu + \partial_\mu f \quad (11)$$

Comparing (11) with (6) we can immediately see that the effect of noncommutativity is precisely that of providing a new symmetry breaking term to the gauge field, instead of the ϕ term, (Cf.refs. [6, 7]) a term not belonging to the gauge field itself.

On the other hand if we neglect in (9) terms $\sim l^2$, then there is no extra contribution coming from (10) or (11), so that we are in the usual non-Abelian gauge field theory, requiring a broken symmetry to obtain an equation like (11).

To see this in greater detail, we note that, as shown by the author in early 2000 [8] given a minimum length l , the energy momentum relation gets modified. The usual Quantum Mechanical commutation relations get modified and now become

$$[x, p] = \hbar' = \hbar \left[1 + \left(\frac{l}{\hbar} \right)^2 p^2 \right] \text{ etc} \quad (12)$$

(Cf. also ref.[9]). (12) shows that effectively $\hbar$ is replaced by $\hbar'$. So, in units, $\hbar = 1 = c$,

$$E = [m^2 + p^2(1 + l^2 p^2)^{-2}]^{\frac{1}{2}}$$

or, the energy-momentum relation leading to the Klein-Gordon Hamiltonian is given by,

$$E^2 = m^2 + p^2 - 2l^2 p^4, \quad (13)$$

neglecting higher order terms. This is the so called Snyder-Sidharth Hamiltonian for Bosons [10]. (It may be mentioned that some other authors have since ad hoc taken a third power of p , and so on [11]. However we should remember that these were mostly phenomenological approaches.)

For Fermions the analysis can be more detailed, in terms of Wilson lattices [12]. The free Hamiltonian now describes a collection of harmonic fermionic oscillators in momentum space. Assuming periodic boundary conditions in all three directions of a cube of dimension L^3 , the allowed momentum components are

$$\mathbf{q} \equiv \left\{ q_k = \frac{2\pi}{L} v_k; k = 1, 2, 3 \right\}, \quad 0 \leq v_k \leq L - 1 \quad (14)$$

(14) finally leads to

$$E_{\mathbf{q}} = \pm \left(m^2 + \sum_{k=1}^3 a^{-2} \sin^2 q_k \right)^{1/2} \quad (15)$$

where $a = l$ is the length of the lattice, this being the desired result leading to

$$E^2 = p^2 e^2 + m^2 c^4 + \alpha l^2 p^4 \quad (16)$$

(16) shows that α is positive, that is for Fermions the Snyder-Sidharth Hamiltonian is given by (16).

2 A Modified Dirac Equation

Once we consider a discrete spacetime structure, the energy momentum relation, as noted, gets modified [13, 12] and we have in units $c = 1 = \hbar$,

$$E^2 - p^2 - m^2 - l^2 p^4 = 0 \quad (17)$$

l being the Planck length. Let us now consider the Dirac equation

$$\{\gamma^\mu p_\mu - m\} \psi \equiv \{\gamma^\circ p^\circ + \Gamma\} \psi = 0 \quad (18)$$

If we include the extra effect shown in (17) we get

$$\left(\gamma^\circ p^\circ + \Gamma + \beta l p^2 \right) \psi = 0 \quad (19)$$

β being a suitable matrix.

Multiplying (19) by the operator

$$\left(\gamma^\circ p^\circ - \Gamma - \beta l p^2 \right)$$

on the left we get

$$p_0^2 - \left(\Gamma \Gamma + \{\Gamma \beta + \beta \Gamma\} + \beta^2 l^2 p^4 \right) \psi = 0 \quad (20)$$

If (20), as in the usual theory, has to represent (17), then we require that the matrix β satisfy

$$\Gamma \beta + \beta \Gamma = 0, \quad \beta^2 = 1 \quad (21)$$

It follows that,

$$\beta = \gamma^5 \quad (22)$$

Using (22) in (19), the modified Dirac equation the so called Dirac-Sidharth equation [10] finally becomes

$$\{\gamma^\circ p^\circ + \Gamma + \gamma^5 l p^2\} \psi = 0 \quad (23)$$

Owing to the fact that we have [14]

$$P\gamma^5 = -\gamma^5 P \quad (24)$$

It follows that the modified Dirac equation (23) is not invariant under reflections.

We can also see that due to the modified Dirac equation (23), there is no additional effect on the anomalous gyromagnetic ratio. This is because, in the usual equation from which the magnetic moment is determined [15] viz.,

$$\frac{d\vec{S}}{dt} = -\frac{e}{\mu c} \vec{B} \times \vec{S},$$

where $\vec{S} = \hbar \Sigma / 2$ is the electron spin operator, there is now an extra term

$$[\gamma^5, \Sigma] \quad (25)$$

However the expression (25) vanishes by the property of the Dirac matrices. We point out that using the modified dispersion relation (16), for a massless particle, $m = 0$, and identifying the extra term $l^2 p^4$ as being due to a mass δm , we can easily deduce that, restoring proper units,

$$\frac{c^2}{\hbar^2} l^2 p^4 = \Delta E^2 = \delta m^2 c^4,$$

Whence,

$$\delta m = \frac{\hbar}{cl} \quad \text{or} \quad l = \frac{\hbar}{c\delta m} \quad (26)$$

This shows that l is the Compton wavelength for this mass δm or alternatively if l is the Compton wavelength, then we deduce the mass, now generated from the extra effect. This is another demonstration of mass generation from $O(l^2)$ effects as seen in Section 1, without requiring a Higgs mechanism. If, for example, l were the Planck length, then δm would be the Planck mass (and vice versa).

3 Remarks

Finally, we would like to make two remarks. Glinka has used the Snyder-Sidharth Hamiltonian (13), to show that a consistent structure is [16] ,

$$E = \frac{p^2}{2m} + m \quad (c = 1) \quad (27)$$

This semi-relativistic solution too, brings out the extra mass (or energy) term.

There is another way of looking at all this. Because of Quantum Theory, the relativistic formula,

$$(\Delta x)^2 - (\Delta t)^2 = 0$$

goes over to, as is well known,

$$(\Delta x)^2 - (\Delta t)^2 \leq \frac{1}{m^2} \quad (\hbar = 1) \quad (28)$$

Effectively, the velocity of light $c (= 1)$ becomes c' , given by from (28),

$$c'^2 \leq 1 + \frac{1}{m^2 \Delta^2} = 2 \quad (29)$$

as $\Delta t \geq \frac{1}{m}$, the Compton time.

So, the usual energy $E = mc^2$ becomes E' where

$$E < E' \leq 2E. \quad (30)$$

There is thus an increase in energy in the extreme relativistic case. This conclusion (30) can also be deduced from (27) if we use (29). Then we get back (30).

Secondly, if we started out with a massless particle like a photon, then using the fact that $E = p = m(c = 1)$, we get directly from (13),

$$l \sim \frac{1}{m}$$

This is in agreement with earlier results [17], giving the photon a mass of $\sim 10^{-65} gm$ and $l \sim$ radius of the universe $\sim 10^{28} cm$.

References

- [1] Sidharth, B.G. (2005). *The Universe of Fluctuations* (Springer, Netherlands).
- [2] Moriyasu, K. (1983). *An Elementary Primer for Gauge Theory* (World Scientific, Singapore).
- [3] Jacob, M. (1974). *Physics Reports, Reprint Volume* (North-Holland, Amsterdam).
- [4] Greiner, W. and Reinhardt, I. (1995). *Gauge Theory of Weak Interactions* (Springer-Verlag, Berlin).
- [5] Taylor, J.C. (1978). *Gauge Theories of Weak Interactions* (Cambridge University Press, Cambridge).
- [6] Sidharth, B.G. (2004). *Proceedings of the Fifth International Symposium on “Frontiers of Fundamental Physics”* (Universities Press, Hyderabad).
- [7] Sidharth, B.G. (2006). *Int.J.SCI.* January 2006, pp.42ff.
- [8] Sidharth, B.G. (2000). *Chaos, Solitons and Fractals* 11, 2000.
- [9] Sidharth, B.G. (2005). *Chaos, Solitons and Fractals* 25, pp.965–968.
- [10] Glinka, L. (2008). *arXiv:hep-th 0812.0551v1*.
- [11] Myers, R.C. and Pospelov, M. (2003). *Phys.Rev.Lett.* 90:211601.
- [12] Montvay, I. and Munster, G. (1994). *Quantum Fields on a Lattice* (Cambridge University Press) pp.174ff.
- [13] Sidharth, B.G. (2005). *Int.J.Mod.Phys.E.* 14, (6), pp.923ff.
- [14] Bjorken, J.D. and Drell, S.D. (1965). *Relativistic Quantum Fields* (McGraw-Hill Inc., New York), pp.44ff.
- [15] E. Merzbacher. (1970). *Quantum Mechanics* (Wiley, New York); Greiner, W. (1983). *Relativistic Quantum Mechanics: Wave Equation* 2nd Ed. (Springer); Greiner, W. and Reinhardt, J. (1987). *Quantum Electrodynamics* 3rd Ed. (Springer).

- [16] Glinka, L.A. (2009). *0811.0551* and *Apeiron* 2, April 2009.
- [17] Sidharth, B.G. (2008) *The Thermodynamic Universe* (World Scientific), Singapore.